

## Appendix F References

1. Renal Association.  
Treatment of adult patients with renal failure. Recommended standards and audit measures.  
Renal Association 1998.
2. Reinsch, C.H. (1967),  
Smoothing by spline functions  
Numerische Mathematik 10,177-183
3. Roderick PJ, Ferris G, Feest TG.  
The provision of renal replacement therapy for adults in England and Wales: recent trends and future directions  
Quarterly Journal of Medicine 1998; 91: 581-587.
4. Department of Health.  
Review of Renal services in England 1993-4  
London: NHS Executive 1996.
5. United States Renal Data System.  
USRDS 1997 Annual Data Report. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD. April 1997.
6. Held PJ, Port FK, Webb RL, et al.  
The USRDS 1997 Annual Data Report  
American Journal of Kidney Diseases 1997; 30; S40-S54.
7. ANZDATA Registry Report 1997  
Australia and New Zealand Dialysis and Transplant Registry, Adelaide, South Australia
8. McDonald JA for the Scottish Renal Registry.  
The Scottish Renal Registry  
Nephrology Dialysis and Transplantation 1998; 13;A121
9. Junor BJR.  
Audit of quality of hospital haemodialysis in Scotland.  
Nephrology Dialysis and Transplantation 1997; 12; 29-32.
10. Berthoux F.  
Annual report on the management of renal failure in Europe XXVIII, 1997.  
Part 1: epidemiological data of treated end stage renal failure in the European Community during the year 1995.  
Presented to the XXXV<sup>th</sup> congress of the EDTA, Rimini, June 1997
11. Lowrie EG, Lowrie MD, Lew NL SM.  
Death Risk in Haemodialysis Patients: The Predictive Value of Commonly Measured Variables and an Evaluation of Death Rate Differences Between Facilities.  
American Journal of Kidney Disease 1990; 15: 5; 458-482.
12. Cavill I.  
Clinics in Haematology  
1982 Vol 11 No 2. 259 - 273